

Bilinear forms on exact operator spaces

and

$$B(H) \otimes B(H).$$

by

Marius Junge and Gilles Pisier*

Abstract.

Let E, F be exact operator spaces (for example subspaces of the C^* -algebra $K(H)$ of all the compact operators on an infinite dimensional Hilbert space H). We study a class of bounded linear maps $u: E \rightarrow F^*$ which we call tracially bounded. In particular, we prove that every completely bounded (in short *c.b.*) map $u: E \rightarrow F^*$ factors boundedly through a Hilbert space. This is used to show that the set OS_n of all n -dimensional operator spaces equipped with the *c.b.* version of the Banach Mazur distance is not separable if $n > 2$.

As an application we show that there is more than one C^* -norm on $B(H) \otimes B(H)$, or equivalently that

$$B(H) \otimes_{\min} B(H) \neq B(H) \otimes_{\max} B(H),$$

which answers a long standing open question.

Finally we show that every “maximal” operator space (in the sense of Blecher-Paulsen) is not exact in the infinite dimensional case, and in the finite dimensional case, we give a lower bound for the “exactness constant”.

Plan.

0. Introduction and background.
1. Factorization of bilinear forms on exact operator spaces.
2. The non-separability of OS_n .
3. Applications to $B(H) \otimes B(H)$ and maximal operator spaces.
4. A new tensor product for C^* -algebras or operator spaces.

* Partially supported by the NSF

§0. Introduction and background.

Following the remarkable results of Kirchberg on exact C^* -algebras *cf.* [Ki1, Ki2], the notion of “exact operator space” was studied in the paper [P1]. In this paper we continue the investigation started in [P1].

Let E, F be operator spaces, we denote

$$d_{cb}(E, F) = \inf\{\|u\|_{cb} \|u^{-1}\|_{cb}\}$$

where the infimum runs over all possible isomorphisms $u: E \rightarrow F$. If E, F are not isomorphic, we set, by convention, $d_{cb}(E, F) = \infty$. We denote by $B(H)$ (resp. $K(H)$) the algebra of all bounded (resp. compact) operators on a Hilbert space H . (See below for unexplained notation and terminology.)

In [P1], the following characteristic of an operator space, E with $\dim E = n$, was studied:

$$(0.1) \quad d_{SK}(E) = \inf\{d_{cb}(E, F) \mid F \subset K(\ell_2)\},$$

or equivalently

$$(0.1)' \quad d_{SK}(E) = \inf\{d_{cb}(E, F) \mid F \subset M_N, N \geq n\}.$$

For an infinite dimensional operator space X , we define

$$d_{SK}(X) = \sup\{d_{SK}(E)\}$$

where the supremum runs over all possible finite dimensional subspaces $E \subset X$.

We say that X is exact if $d_{SK}(X) < \infty$. This is equivalent (see [P1]) to the exactness of a certain sequence of morphisms in the category of operator spaces, whence the terminology.

Let E, F be exact operator spaces. We will prove below an inequality for completely bounded linear maps $u: E \rightarrow F^*$. Here F^* means the “operator space dual” of F in the sense of [ER1, BP].

This key inequality can be viewed as a form of Grothendieck’s inequality for exact operator spaces. It implies that every *c.b.* map $u: E \rightarrow F^*$ can be factorized (in the Banach

space sense) through a Hilbert space with a norm of factorization $\leq 4\|u\|_{cb}$. Actually, in this inequality “complete boundedness” can be replaced by a more general notion which we call “tracial boundedness” which has already been considered by previous authors (*cf.* [I, B3]). Using this key inequality and a somewhat surprising application of Baire’s theorem, we prove that the metric space OS_n of all n -dimensional operator spaces equipped with the distance

$$\delta_{cb}(E, F) = \log d_{cb}(E, F)$$

is not separable as soon as $n > 2$. We can even show that the subset of all isometrically Hilbertian and homogeneous (in the sense of [P3]) n -dimensional operator spaces, is non-separable.

This has a surprising application to C^* -algebra theory. Recall that a C^* -algebra A is called nuclear if $A \otimes_{\min} B = A \otimes_{\max} B$ for any C^* -algebra B (see [La]). For a long time it remained an open problem whether it suffices for the nuclearity of A to assume that

$$(*) \quad A \otimes_{\min} A^{op} = A \otimes_{\max} A^{op}$$

where A^{op} is the opposite C^* -algebra (with the product in reverse order). In [Ki2], Kirchberg gave the first counterexamples. However, he pointed out that it remained unknown whether $(*)$ holds in the (non-nuclear) case of $A = B(H)$. (Note that $B(H)$ is isomorphic to its opposite.) Kirchberg also proposed an approach to this question together with a series of equivalent conjectures. One of his conjectures was our main motivation to investigate the non-separability of the metric space OS_n , and as a result, we obtain a negative answer to the above mentioned question: the identity $(*)$ fails for $A = B(H)$. In other words, we have

$$B(H) \otimes_{\min} B(H) \neq B(H) \otimes_{\max} B(H).$$

Equivalently, there is more than one C^* -norm on $B(H) \otimes B(H)$ whenever H is an infinite dimensional Hilbert space. This is proved in section 3.

In the same section, we also show that if E is an n -dimensional Banach space then the operator space $\max(E)$ in the sense of [BP] satisfies

$$d_{SK}(\max(E)) \geq \frac{\sqrt{n}}{4}.$$

In particular, for any infinite dimensional Banach space X the operator space $\max(X)$ is not exact.

After circulating a first version of this paper, we observed that the non-separability of OS_n for $n > 2$ can be alternately deduced from properties of Kazhdan-groups, following the ideas of Voiculescu [V] (see Remark 2.10 below). However, this approach does not seem to give quite as much information as our original one.

In the final section 4, we introduce a "new" tensor product obtained as follows. Let $E \subset B(H)$ and $F \subset B(K)$ be two operator spaces, we denote by $E \otimes_M F$ the completion of the linear tensor product with respect to the norm induced on it by $B(H) \otimes_{\max} B(K)$. This tensor product makes sense both in the category of operator spaces and in that of C^* -algebras. Our previous results show that it differs in general from the minimal tensor product. We include several properties of this tensor product, based mainly on [Ki3].

In the rest of this introduction we give some background and explain our notation. We refer to [Sa, Ta] for operator algebra theory. Let H, K be Hilbert spaces. We will denote by $H \otimes_2 K$ their Hilbertian tensor product. By an operator space we mean a closed subspace of $B(H)$ for some Hilbert space H . If $E_1 \subset B(H_1)$, $E_2 \subset B(H_2)$ are operator spaces, we will denote by $E_1 \otimes_{\min} E_2$ their minimal (or spatial) tensor product equipped with the minimal (or spatial) tensor norm induced by the space $B(H_1 \otimes_2 H_2)$. When E_1 and E_2 are C^* -subalgebras then this norm is a C^* -norm, and actually it is the smallest C^* -norm on the algebraic tensor product $E_1 \otimes E_2$. In the case of C^* -algebras, we will denote by $E_1 \otimes_{\max} E_2$ the completion of $E_1 \otimes E_2$ with respect to the largest C^* -norm on $E_1 \otimes E_2$.

We recall that if $E \subset B(H)$ and $F \subset B(K)$ are operator spaces, then a map $u: E \rightarrow F$ is completely bounded (in short *c.b.*) if the maps $u_m = I_{M_m} \otimes u: M_m(E) \rightarrow M_m(F)$ are uniformly bounded when $m \rightarrow \infty$, i.e. if we have $\sup_{m \geq 1} \|u_m\| < \infty$. The *c.b.* norm of u is defined as

$$\|u\|_{cb} = \sup_{m \geq 1} \|u_m\|.$$

We will say that u is completely isometric (resp. completely contractive) or is a complete isometry (resp. a complete contraction) if the maps u_m are isometries (resp. of norm ≤ 1)

for all m . This is the same as saying that for any operator space X the map

$$I_X \otimes u: X \otimes_{\min} E \rightarrow X \otimes_{\min} F$$

is an isometry (resp. a contraction). We also recall that u is called completely positive (in short *c.p.*) if all the maps u_m are positive.

If E_1, F_1, E_2, F_2 are operator spaces and if

$$u_1: E_1 \rightarrow F_1 \quad u_2: E_2 \rightarrow F_2$$

are completely bounded, then

$$u_1 \otimes u_2: E_1 \otimes_{\min} F_1 \longrightarrow E_2 \otimes_{\min} F_2$$

is *c.b.* and we have

$$(0.2) \quad \|u_1 \otimes u_2\|_{cb} = \|u_1\|_{cb} \|u_2\|_{cb}.$$

If $u: E_1 \rightarrow F_1$ and $u_2: E_2 \rightarrow F_2$ are completely isometric then $u_1 \otimes u_2: E_1 \otimes_{\min} E_2 \longrightarrow F_1 \otimes_{\min} F_2$ also is completely isometric. In particular, we note the completely isometric identity

$$M_n(E) = M_n \otimes_{\min} E.$$

We will use (0.2) repeatedly in the sequel with no further reference. We refer the reader to [Pa1] for more information.

It is known that the analogue of (0.2) fails for the max-tensor product. Instead we have:

(0.3) If E_1, F_1 and E_2, F_2 are C^* -algebras (or merely operator systems) and if $u_1: E_1 \rightarrow F_1$ and $u_2: E_2 \rightarrow F_2$ are completely positive maps, then the map $u_1 \otimes u_2$ defined on the algebraic tensor product extends to a completely positive and *c.b.* map

$$u_1 \otimes u_2: E_1 \otimes_{\max} F_1 \longrightarrow E_2 \otimes_{\max} F_2$$

satisfying

$$\|u_1 \otimes u_2\|_{cb(E_1 \otimes_{\max} F_1, E_2 \otimes_{\max} F_2)} = \|u_1\| \|u_2\|.$$

See e.g. [Ta, Proposition 4.23], [Pa1, Proposition 3.5 and Proposition 10.11] or [W2, Proposition 1.11] for more details.

We will use the duality theory for operator spaces, which was introduced in [BP, ER1] using Ruan's "abstract" characterization of operator spaces ([Ru]). This can be summarized as follows: Let E be an operator space and let E^* be the dual Banach space. Then, for some Hilbert space K , there is an isometric embedding $E^* \subset B(K)$ such that for any operator space F , the minimal (=spatial) norm on $E^* \otimes F$ coincides with the norm induced on it by the space $cb(E, F)$. Moreover, this property characterizes the operator space

$$E^* \subset B(K)$$

up to complete isometry. We will refer to this operator space as the "operator space dual" of E . We refer to [BP, ER1-2, B1-2, Ru] for detailed information.

Consider arbitrary Banach spaces E, Y and an operator $u: E \rightarrow Y$.

Recall that u is called 2-absolutely summing if there is a constant C such that for all n for all $(x_1, \dots, x_n)$ in E we have

$$\sum \|ux_i\|^2 \leq C^2 \sup \left\{ \sum |\xi(x_i)|^2 \mid \xi \in E^*, \|\xi\| \leq 1 \right\}.$$

We denote by $\pi_2(u)$ the smallest constant C for which this holds.

It is easy to check that for any bounded operators $v: Y \rightarrow Y_1$ and $w: E_1 \rightarrow E$ we have

$$(0.4) \quad \pi_2(vuw) \leq \|v\| \pi_2(u) \|w\|.$$

Notation: Let (x_i) be a finite sequence in a C^* -algebra A we will denote for brevity

$$RC((x_i)) = \max \left\{ \left\| \sum x_i x_i^* \right\|, \left\| \sum x_i^* x_i \right\| \right\}.$$

Note that if A is commutative and if $E \subset A$ is any subspace equipped with the induced norm, we have for all $(x_1, \dots, x_n)$ in E

$$(0.5) \quad RC((x_i)) = \sup \left\{ \sum |\xi(x_i)|^2 \mid \xi \in E^*, \|\xi\| \leq 1 \right\}.$$

Definition 0.3. Consider a C^* -algebra A and a Banach space Y . Let $E \subset A$ be subspace. A linear map $u: E \rightarrow Y$ will be called $(2, RC)$ -summing if there is a constant C such that for all n and for all $(x_1, \dots, x_n)$ in E we have

$$\sum \|ux_i\|^2 \leq C^2 RC((x_i)).$$

We denote by $\pi_{2,RC}(u)$ the smallest constant C for which this holds. We refer to [P3] for a more systematic treatment of $(2, w)$ -summing operators when w is a “weight” in the sense of [P3]. (See also [P4, section 2] or [P2].) By a well known variant of a Pietsch’s factorization theorem for 2-absolutely summing operators, we have

Proposition 0.4. Consider $u: E \rightarrow Y$ as in the preceding definition. The following are equivalent.

- (i) u is $(2, RC)$ -summing and $\pi_{2,RC}(u) \leq C$.
- (ii) There are states f, g on A and $0 \leq \theta \leq 1$ such that

$$\forall x \in E \quad \|u(x)\| \leq C\{\theta f(x^*x) + (1 - \theta)g(xx^*)\}^{1/2}.$$

- (iii) The map $u: E \rightarrow Y$ admits an extension $\tilde{u}: A \rightarrow Y$ such that $\pi_{2,RC}(\tilde{u}) \leq C$.

Proof. This is – by now – a well known application of the Hahn-Banach theorem. For more details we refer the reader e.g. to [P4, Lemma 1.3] or [P3, Prop. 5.1].

For bilinear forms, we have the following known analogous statement. (In the commutative case, this can be found in [Kw].)

Proposition 0.5. Let A, B be C^* -algebras and let $E \subset A, F \subset B$ be closed subspaces. Let $C > 0$ be a fixed constant. The following properties of a linear map $u: E \rightarrow F^*$ are equivalent.

- (i) For any n , any $(x_1, \dots, x_n)$ in E and any $(y_1, \dots, y_n)$ in F , we have

$$\left| \sum \langle u(x_i), y_i \rangle \right| \leq C[RC((x_i))RC((y_i))]^{1/2}.$$

- (ii) There are states f_1, g_1 on A, f_2, g_2 on B and $0 \leq \theta_1, \theta_2 \leq 1$ such that

$$\begin{aligned} \forall (x, y) \in E \times F \quad |\langle u(x), y \rangle| &\leq C[\theta_1 f_1(x^*x) + (1 - \theta_1)g_1(xx^*)]^{1/2} \\ &\quad [\theta_2 f_2(y^*y) + (1 - \theta_2)g_2(yy^*)]^{1/2}. \end{aligned}$$

(iii) For some Hilbert space H , u admits a factorization of the form

$$E \xrightarrow{a} H \xrightarrow{b^*} F^*$$

with operators $a: E \rightarrow H$ and $b: F \rightarrow H^*$ such that

$$\pi_{2,RC}(b)\pi_{2,RC}(a) \leq C.$$

Proof. See e.g. [P3, Theorem 6.1] or [P4, Lemma 1.3].

Notation: Let $u: X \rightarrow Y$ be an operator between Banach spaces. Assume that u factors through a Hilbert space H , i.e. we have $u = \alpha\beta$ with $\alpha: H \rightarrow Y$ and $\beta: X \rightarrow H$. Then we will denote

$$\gamma_2(u) = \inf\{\|\alpha\| \|\beta\|\}$$

where the infimum runs over all possible factorization. This is the “norm of factorization through Hilbert space” of u . See [P2] for more on this theme.

Corollary 0.6. *Let u be as in Proposition 0.5. Then if (iii) holds there are operators $\tilde{a}: A \rightarrow H$ and $\tilde{b}: B \rightarrow H^*$ such that $\tilde{b}^*\tilde{a}$ (viewed as a bilinear form on $A \times B$) extends u and*

$$\pi_{2,RC}(\tilde{a})\pi_{2,RC}(\tilde{b}) \leq C.$$

In particular, if we let $\tilde{u} = \tilde{b}^\tilde{a}$, then the operator $\tilde{u}: A \rightarrow B^*$ satisfies $\|\tilde{u}\| \leq C$ and $\langle \tilde{u}(x), y \rangle = \langle u(x), y \rangle$ for all (x, y) in $E \times F$. Moreover we have $\gamma_2(u) \leq C$.*

Proof. This follows from Propositions 0.4 and 0.5.

The following fact is well known to specialists. (See e.g. [Ki1] Lemma 3.9)

Lemma 0.7. *Let C be a C^* -algebra and let $I \subset C$ be a closed ideal. Let E, X be operator spaces with $E \subset X$. Consider the canonical (complete) contraction $C \otimes_{\min} X \rightarrow (C/I) \otimes_{\min} X$. Since this map vanishes on $I \otimes_{\min} X$, we clearly have a canonical (complete) contraction*

$$T_X: C \otimes_{\min} X / I \otimes_{\min} X \rightarrow (C/I) \otimes_{\min} X.$$

If T_X is an isomorphism, then T_E also is an isomorphism, moreover

$$\|T_E^{-1}\| \leq \|T_X^{-1}\|.$$

Proof. It is well known (cf. [Ta, p. 27]) that I possesses an approximate unit formed of elements $p_i \in I$ such that $0 \leq p_i \leq 1$ (hence $\|1 - p_i\| \leq 1$) and $p_i x \rightarrow x \forall x \in I$. Then the proof can be completed exactly as in [P1, Lemma 3].

Remark. Equivalently, if we consider the complete isometry

$$C \otimes_{\min} E \rightarrow C \otimes_{\min} X$$

then this map defines after passing to the quotient spaces a complete isometry

$$C \otimes_{\min} E / I \otimes_{\min} E \rightarrow C \otimes_{\min} X / I \otimes_{\min} X.$$

We will also invoke the following elementary fact which follows easily (like the preceding result) from the existence of an approximate unit in any ideal of a C^* -algebra. (Recall that $A \otimes B$ denotes the algebraic tensor product.)

Lemma 0.8. *Let A, B, C be C^* -algebras. Let $\pi: C \rightarrow A$ be a surjective $*$ -homomorphism. Let $\mathcal{I} = \text{Ker}(\pi)$. Then (viewing the three sets appearing below as subsets of $C \otimes_{\min} B$) we have*

$$[C \otimes B] \cap [\mathcal{I} \otimes_{\min} B] = \mathcal{I} \otimes B.$$

Equivalently, let

$$q: C \otimes_{\min} B \rightarrow [C \otimes_{\min} B] / [\mathcal{I} \otimes_{\min} B]$$

be the quotient map and let

$$T: [C \otimes_{\min} B] / [\mathcal{I} \otimes_{\min} B] \rightarrow A \otimes_{\min} B$$

be the morphism associated to $\pi \otimes I_B$. Then T induces a linear and $*$ -algebraic isomorphism between $q(C \otimes B)$ and $A \otimes B$.

Acknowledgement: The second author would like to thank E. Kirchberg for introducing him to the questions considered in this paper and B. Maurey for a stimulating conversation.

§1. Factorization of bilinear forms on exact operator spaces.

Let $n \geq 1$ be an integer. We will denote by

$$J_n: M_n \rightarrow M_n^*$$

the map defined by

$$\forall x, y \in M_n \quad \langle J_n(x), y \rangle = \frac{1}{n} \text{tr}(^t y x).$$

The following notion is natural for our subsequent results. It has already been considered in [I].

Definition 1.1. *Let E, F be operator spaces. Let $u: E \rightarrow F^*$ be a linear map. We will say that u is tracially bounded (in short t.b.) if*

$$\sup_{n \geq 1} \|J_n \otimes u\|_{M_n(E) \rightarrow M_n(F)^*} < \infty$$

and we denote

$$\|u\|_{tb} = \sup_{n \geq 1} \|J_n \otimes u\|_{M_n(E) \rightarrow M_n(F)^*}.$$

Equivalently, $u: E \rightarrow F^*$ is t.b. iff the bilinear forms

$$u_n: M_n(E) \times M_n(F) \rightarrow \mathbf{C}$$

defined by

$$u_n((x_{ij}), (y_{ij})) = \frac{1}{n} \sum_{ij} \langle u(x_{ij}), y_{ij} \rangle$$

are bounded uniformly in n and we have

$$\|u\|_{tb} = \sup_{n \geq 1} \|u_n\|.$$

We immediately observe

Lemma 1.2. *For a linear map $u: E \rightarrow F^*$*

$$\text{complete boundedness} \Rightarrow \text{tracial boundedness} \Rightarrow \text{boundedness},$$

and we have

$$\|u\| \leq \|u\|_{tb} \leq \|u\|_{cb}.$$

Proof. If u is *c.b.* then for any u and any $(x_{ij}) \in M_n(E)$

$$\|(u(x_{ij}))\|_{M_n(F^*)} \leq \|u\|_{cb} \|x\|_{M_n(E)}.$$

We have $M_n(F^*) = cb(F, M_n)$, hence

$$\begin{aligned} \|(u(x_{ij}))\|_{M_n(F^*)} &= \sup\{\|\langle u(x_{ij}), y_{k\ell} \rangle\|_{M_n(M_n)} \mid \|(y_{k\ell})\|_{M_n(F)} \leq 1\} \\ &= \sup_{y \in B_{M_n(F)}} \left\{ \left| \sum_{ijk\ell} \langle u(x_{ij}), y_{k\ell} \rangle \alpha_{j\ell} \beta_{ik} \right| \mid \sum_{j\ell} |\alpha_{j\ell}|^2 \leq 1, \sum_{ik} |\beta_{ik}|^2 \leq 1 \right\}. \end{aligned}$$

Taking $\alpha, \beta = \frac{I}{\sqrt{n}}$ we get

$$\|(u(x_{ij}))\|_{M_n(F^*)} \geq \sup \left\{ \frac{1}{n} \left| \sum \langle u(x_{ij}), y_{ij} \rangle \right| \mid \|(y_{ij})\|_{M_n(F)} \leq 1 \right\}$$

hence $\|u\|_{tb} \leq \|u\|_{cb}$. The inequality $\|u\| \leq \|u\|_{tb}$ is clear by taking $n = 1$. ■

The following consequence of the non-commutative Grothendieck inequality is known ([B3]).

Lemma 1.3. *Let A, B be C^* -algebras. Then any bounded linear operator $u: A \rightarrow B^*$ is tracially bounded and we have*

$$\|u\| \leq \|u\|_{tb} \leq K \|u\|$$

for some numerical constant K . Let $E \subset A$ and $F \subset B$ be closed subspaces and let $i_E: E \rightarrow A$, $i_F: F \rightarrow B$ be the inclusion maps. Then the restriction $i_F^* u i_E: E \rightarrow F^*$ satisfies

$$\|i_F^* u i_E\|_{tb} \leq K \|u\|.$$

Proof. Consider $x = (x_{ij}) \in M_n(A)$ and $y = (y_{ij}) \in M_n(B)$.

Let $\alpha(x) = \max \left\{ n^{-1/2} \left\| \sum_{ij} x_{ij}^* x_{ij} \right\|^{1/2}, n^{-1/2} \left\| \sum_{ij} x_{ij} x_{ij}^* \right\|^{1/2} \right\}$. It is easy to check that

$$\alpha(x) \leq \|x\|_{M_n(A)}.$$

By the non-commutative Grothendieck inequality (cf. [H1], see also [P2]) we have for some numerical absolute constant K

$$\left| \frac{1}{n} \sum_{ij} \langle u(x_{ij}), y_{ij} \rangle \right| \leq K \|u\| \alpha(x) \alpha(y)$$

hence a fortiori

$$\leq K \|u\| \|x\|_{M_n(A)} \|y\|_{M_n(B)}.$$

Therefore $\|u\|_{tb} \leq K \|u\|$. The second part is obvious since

$$\|i_F^* u i_E\|_{tb} \leq \|u\|_{tb}.$$

■

Remark. The preceding argument actually shows the following: Consider $u: A \rightarrow B^*$ of the form $u = a^*b$, where $b: E \rightarrow H$ and $a: F \rightarrow H^*$ are $(2, RC)$ -summing operators. Then we have

$$\|u\|_{tb} \leq \pi_{2,RC}(a) \pi_{2,RC}(b).$$

We will now show that if E and F are exact operator spaces, the converse to the second part of Lemma 1.3 also holds, that is to say the bilinear form on $E \times F$ associated to a tracially bounded map $u: E \rightarrow F^*$ is the restriction of a bounded bilinear form on $A \times B$. This is the key result for this paper.

We denote by F_∞ the free group with countably many generators denoted by $g_1, g_2, \dots$. Let $\lambda: F_\infty \rightarrow B(\ell_2(F_\infty))$ be the left regular representation and let us denote simply by C_λ the reduced C^* -algebra of F_∞ . Let E be an operator space. Let $(x_t)_{t \in F_\infty}$ be a finitely supported family of elements of E . For simplicity we will denote simply by

$$\left\| \sum \lambda(t) \otimes x_t \right\|_{\min}$$

the norm of $\sum_{t \in F_\infty} \lambda(t) \otimes x_t$ in $C_\lambda \otimes_{\min} E$.

The following inequality (cf. [HP, Prop. 1.1]) plays an important rôle in the sequel.

(1.1) For any finite sequence (x_i) in a C^* -algebra we have

$$\left\| \sum \lambda(g_i) \otimes x_i \right\|_{\min} \leq 2 \max \left\{ \left\| \sum x_i^* x_i \right\|^{1/2}, \left\| \sum x_i x_i^* \right\|^{1/2} \right\}.$$

Theorem 1.4. *Let E, F be exact operator spaces. Let $C = d_{SK}(E)d_{SK}(F)$. Let $u: E \rightarrow F^*$ be a tracially bounded linear map. Let $(x_t)_{t \in F_\infty}$ (resp. $(y_t)_{t \in F_\infty}$) be a finitely supported family of elements of E (resp. F). Then we have*

$$(1.2) \quad \left| \sum \langle u(x_t), y_t \rangle \right| \leq C \|u\|_{tb} \left\| \sum \lambda(t) \otimes x_t \right\|_{\min} \left\| \sum \lambda(t) \otimes y_t \right\|_{\min},$$

In particular, for all n and all $x_i \in E, y_i \in F$ ($i = 1, 2, \dots, n$) we have

$$(1.3) \quad \left| \sum \langle u(x_i), y_i \rangle \right| \leq C \|u\|_{tb} \left\| \sum \lambda(g_i) \otimes x_i \right\|_{\min} \left\| \sum \lambda(g_i) \otimes y_i \right\|_{\min},$$

hence

$$(1.3)' \quad \leq 4C \|u\|_{tb} \max \left\{ \left\| \sum x_i^* x_i \right\|^{1/2}, \left\| \sum x_i x_i^* \right\|^{1/2} \right\} \max \left\{ \left\| \sum y_i^* y_i \right\|^{1/2}, \left\| \sum y_i y_i^* \right\|^{1/2} \right\}.$$

Furthermore, if A, B are C^ -algebras such that $E \subset A, F \subset B$ (completely isometrically) then u admits an extension $\tilde{u}: A \rightarrow B^*$ such that*

$$\|\tilde{u}\| \leq \|\tilde{u}\|_{tb} \leq 4C \|u\|_{tb}$$

and $\langle \tilde{u}(x), y \rangle = \langle u(x), y \rangle \forall (x, y) \in E \times F$.

Proof. The proof is essentially the same as that of in [P1, Theorem 8]. An essential ingredient in the proof is Wassermann's construction of a specific embedding of C_λ into an ultrapower (in the von Neumann sense) of matrix algebras, cf. [W1]. This is based on the residual finiteness of the free group.

More precisely, consider the family

$$\{M_\alpha \mid \alpha \geq 1\}$$

of all matrix algebras. Let $L = \{(x_\alpha)_{\alpha \geq 1} \mid x_\alpha \in M_\alpha, \sup_{\alpha \geq 1} \|x_\alpha\|_{M_\alpha} < \infty\}$ equipped with the norm $\|(x_\alpha)\|_L = \sup_{\alpha \geq 1} \|x_\alpha\|_{M_\alpha}$. Let $\mathcal{U}$ be an ultrafilter on $\mathbf{N}$. Let us denote by τ_α the normalized trace on M_α , and let

$$I_{\mathcal{U}} = \{(x_\alpha)_{\alpha \geq 1} \in L \mid \lim_{\mathcal{U}} \tau_\alpha(x_\alpha^* x_\alpha) = 0\}.$$

We then set $\mathcal{N} = L/I_{\mathcal{U}}$. It is a well known fact (cf [Sa]) that $\mathcal{N}$ is a finite von Neumann algebra. Let VN_{λ} be the von Neumann algebra generated by λ . (Note that $C_{\lambda} \subset VN_{\lambda}$.) Wassermann proved that for a suitable $\mathcal{U}$ one can find for each g in F_{∞} a sequence $(u_{\alpha}^g)_{\alpha \geq 1}$ such that:

- (i) u_{α}^g is unitary in M_{α} and has real entries ($\alpha \geq 1, i = 1, 2, \dots$),
- (ii) $\lim_{\mathcal{U}} \tau_{\alpha}(u_{\alpha}^{s*} u_{\alpha}^g) = 0$ if $g \neq s$, or equivalently since the entries are real $\lim_{\mathcal{U}} \tau_{\alpha}(^t u_{\alpha}^s u_{\alpha}^g) = 0$ if $g \neq s$.
- (iii) The mapping $\Phi: VN_{\lambda} \rightarrow L/I_{\mathcal{U}}$ which takes $\lambda(g)$ to the equivalence class of $(u_{\alpha}^g)_{\alpha \geq 1}$ is an isometric representation mapping VN_{λ} onto a von Neumann subalgebra of $\mathcal{N} = L/I_{\mathcal{U}}$. A fortiori it is completely isometric.

The last point implies that we can write

$$(1.4) \quad \left\| \sum_{t \in F_{\infty}} \lambda(t) \otimes x_t \right\|_{\min} = \left\| \sum_{t \in F_{\infty}} \Phi[\lambda(t)] \otimes x_t \right\|_{(L/I_{\mathcal{U}}) \otimes_{\min} E}.$$

Without loss of generality we may assume $\dim E = \dim F = n$. Hence, for any $\varepsilon > 0$, for some integer N there is $E_1 \subset M_N$ such that

$$(1.5) \quad d_{cb}(E, E_1) < d_{SK}(E)(1 + \varepsilon).$$

Clearly we have completely isometrically

$$L/I_{\mathcal{U}} \otimes_{\min} M_N = (L \otimes_{\min} M_N)/I_{\mathcal{U}} \otimes_{\min} M_N.$$

By Lemma 0.7, since $I_{\mathcal{U}}$ is an ideal this remains true with E_1 in the place of M_N . By (1.5), it follows that the natural (norm one) map

$$T_E: (L \otimes_{\min} E)/(I_{\mathcal{U}} \otimes_{\min} E) \rightarrow (L/I_{\mathcal{U}}) \otimes_{\min} E$$

has an inverse with norm $\|T_E^{-1}\| < d_{SK}(E)(1 + \varepsilon)$, and since ε is arbitrary and the same is true for F , we actually have

$$(1.6) \quad \|T_E^{-1}\| \leq d_{SK}(E) \quad \|T_F^{-1}\| \leq d_{SK}(F).$$

Since u is tracially bounded, for any α , the linear map $V_\alpha: M_\alpha(E) \rightarrow M_\alpha(F)^*$ defined by

$$\langle V_\alpha(a_\alpha \otimes x), b_\alpha \otimes y \rangle = \tau_\alpha({}^t b_\alpha a_\alpha) \langle u(x), y \rangle, \quad \forall x \in E, \forall y \in F, \forall a_\alpha, b_\alpha \in M_\alpha$$

is bounded with

$$(1.7) \quad \|V_\alpha\| \leq \|u\|_{tb}.$$

For $a \in L$, let us denote by (a_α) its coordinates. Similarly consider an element z in $L \otimes E$. Clearly z can be identified to a family (z_α) with $z_\alpha \in M_\alpha \otimes E$. Note that

$$(1.8) \quad \|z\|_{L \otimes_{\min} E} = \sup_{\alpha} \|z_\alpha\|_{M_\alpha(E)}.$$

With this notation, we can define a linear map $V: L \otimes_{\min} E \rightarrow (L \otimes_{\min} F)^*$ by setting

$$\langle V(a \otimes x), b \otimes y \rangle = \lim_{\mathcal{U}} \tau_\alpha({}^t b_\alpha a_\alpha) \langle u(x), y \rangle \quad \forall a, b \in L, \forall x \in E, \forall y \in F.$$

Clearly, by (1.7) and (1.8) we have $\|V\| \leq \sup_{\alpha} \|V_\alpha\| \leq \|u\|_{tb}$. Moreover, it is clear that for all ξ in $I_{\mathcal{U}} \otimes_{\min} E$ and all η in $I_{\mathcal{U}} \otimes_{\min} F$ we have $\langle V(\xi), \eta \rangle = 0$. Therefore V defines canonically a map

$$\tilde{V}: (L \otimes_{\min} E) / (I_{\mathcal{U}} \otimes_{\min} E) \rightarrow [(L \otimes_{\min} F) / (I_{\mathcal{U}} \otimes_{\min} F)]^*$$

such that $\|\tilde{V}\| = \|V\| \leq \|u\|_{tb}$. By (1.6), $\tilde{V}$ also defines a map

$$\hat{V}: (L/I_{\mathcal{U}}) \otimes_{\min} E \rightarrow [(L/I_{\mathcal{U}}) \otimes_{\min} F]^*$$

such that

$$(1.9) \quad \|\hat{V}\| \leq C\|u\|_{tb}.$$

Let (x_t) and (y_t) be as in Theorem 1.4, and let $T_1 = \sum \lambda(t) \otimes x_t$, $T_2 = \sum \lambda(t) \otimes y_t$, and let

$$\hat{T}_1 = (\Phi \otimes I_E)(T_1) \in (L/I_{\mathcal{U}}) \otimes_{\min} E, \quad \hat{T}_2 = (\Phi \otimes I_F)(T_2) \in (L/I_{\mathcal{U}}) \otimes_{\min} F.$$

By (ii) above, we clearly have $\sum \langle u(x_t), y_t \rangle = \langle \hat{V}(\hat{T}_1), \hat{T}_2 \rangle$ hence

$$|\sum \langle u(x_t), y_t \rangle| \leq \|\hat{V}\| \|\hat{T}_1\|_{\min} \|\hat{T}_2\|_{\min},$$

and this together with (1.4) and (1.9) implies (1.2). This proves (1.2). Clearly (1.3) is but a particular case of (1.2) and (1.3)' follows, using (1.1).

Finally, the last assertion follows from Corollary 0.6 and the remark after Lemma 1.3. ■

Remark. The preceding proof of (1.2) remains valid if we replace F_∞ by any residually finite discrete group G . Actually, we only use the fact that there is a completely isometric embedding into an ultraproduct, say $\Phi: C_\lambda^*(G) \rightarrow L/I_\mathcal{U}$ such that

$$\forall s, g \in G \quad \lim_{\mathcal{U}} \tau_\alpha({}^t \Phi_\alpha(\lambda(s)) \Phi_\alpha(\lambda(g))) = \delta_{sg}.$$

A corollary of the non-commutative Grothendieck theorem says that every bounded linear operator $u: A \rightarrow B^*$ factors through a Hilbert space when A, B are C^* -algebras and we have $\gamma_2(u) \leq K\|u\|$ for some absolute constant K . In the same vein, we have

Corollary 1.5. *Let E, F be as in Theorem 1.4 with $C = d_{SK}(E)d_{SK}(F)$. Then, if $u: E \rightarrow F^*$ is tracially bounded there is a Hilbert space H and a factorization $u = a^*b$*

$$E \xrightarrow{b} H \xrightarrow{a^*} F^*$$

with a, b such that

$$\pi_{2,RC}(a)\pi_{2,RC}(b) \leq 4C\|u\|_{tb}.$$

Conversely, if there is such a factorization we have

$$(1.10) \quad \|u\|_{tb} \leq \pi_{2,RC}(a)\pi_{2,RC}(b).$$

Proof. The direct implication follows from Theorem 1.4 and Proposition 0.5. The converse follows from the remark after Lemma 1.3. ■

Corollary 1.6. *Let E, F be exact operator spaces. Let $C = d_{SK}(E)d_{SK}(F)$. Then every completely bounded map $u: E \rightarrow F^*$ factors through a Hilbert space and we have*

$$(1.11) \quad \gamma_2(u) \leq 4C\|u\|_{cb}.$$

Proof. First recall that $\|u\|_{tb} \leq \|u\|_{cb}$ by Lemma 1.2. Then this is deduced from Theorem 1.4 using Proposition 0.5 and Corollary 0.6. ■

Corollary 1.7. *Let E be a closed subspace of a commutative C^* -algebra and let F be an exact operator space. Then every completely bounded $u: E \rightarrow F^*$ is 2-absolutely summing and we have $\pi_2(u) \leq 4d_{SK}(F)\|u\|$.*

Proof. Since a commutative C^* -algebra is nuclear we have $d_{SK}(E) = 1$. The result then follows from Theorem 1.4 taking (0.5) into account. ■

In particular we have the following corollary which is already known. It was proved independently by V. Paulsen and the second author on one hand (using Clifford matrices, this gives the better constant 2, see [Pa3]) and by the first author on the other (using random matrices, this yields a worse constant).

Corollary 1.8. *Consider C^* -algebras A, B and subspaces $E \subset A$ and $F \subset B$. If A, B are assumed commutative, then any c.b. map $u: E \rightarrow F^*$ can be written as $u = a^*b$ with $\pi_2(a)\pi_2(b) \leq 4\|u\|_{cb}$.*

Proof. Again, by the nuclearity of A, B we have $d_{SK}(E) = d_{SK}(F) = 1$. Note that by (0.5) we have $\pi_2(a) = \pi_{2,RC}(a)$ for all $a: E \rightarrow H$ and similarly for F . Hence this follows from Theorem 1.4. ■

Corollary 1.9. *Let E be an exact operator space. Consider a linear map $v: E \rightarrow H$ into a Hilbert space. Let $C > 0$ be a constant. Assume that for all n and all (x_{ij}) in $M_n(E)$ we have*

$$(1.12) \quad \left(\frac{1}{n} \sum_{ij} \|v(x_{ij})\|^2 \right)^{1/2} \leq C \|(x_{ij})\|_{M_n(E)}.$$

Then v is $(2, RC)$ -summing and

$$(1.13) \quad \pi_{2,RC}(v) \leq 2Cd_{SK}(E).$$

Proof. Consider the mapping $u = \bar{v}^*v: E \rightarrow \bar{E}^*$, obtained by identifying H with its antidual $\bar{H}^*$. From (1.12) it is easy to deduce by Cauchy-Schwarz that $\|\bar{v}^*v\|_{tb} \leq C^2$. Hence by (1.2) we have for all (x_i) in E

$$\sum \|vx_i\|^2 = \sum \langle \bar{v}^*vx_i, x_i \rangle \leq 4C^2 d_{SK}(E)^2 RC((x_i))^2$$

and (1.13) follows. ■

We need to recall some elementary facts on ultraproducts of operator spaces. Let $(E_i)_{i \in I}$ be a family of operator spaces with $E_i \subset B(H_i)$. Then their ultraproduct $\hat{E} = \Pi E_i / \mathcal{U}$ embeds isometrically into $\Pi B(H_i) / \mathcal{U}$. The latter being a C^* -algebra, this embedding defines an operator space structure on $\hat{E}$. It is easy to check that we have isometrically

$$M_n(\hat{E}) = \Pi M_n(E_i) / \mathcal{U}.$$

Equivalently

$$M_n \otimes_{\min} \hat{E} = \Pi(M_n \otimes_{\min} E_i) / \mathcal{U}.$$

This identity clearly remains valid with M_n replaced by any subspace $F \subset M_n$, therefore we also have the following isometric identity, valid if $d_{SK}(F) = 1$ ($\dim F < \infty$).

$$(1.14) \quad F \otimes_{\min} \hat{E} = \Pi(F \otimes_{\min} E_i) / \mathcal{U}.$$

This yields

Corollary 1.10. *Let I be any set. Let $(E_i)_{i \in I}$, $(F_i)_{i \in I}$ be exact operator spaces with*

$$C = \sup_{i \in I} d_{SK}(E_i) d_{SK}(F_i) < \infty.$$

Let $u_i: E_i \rightarrow F_i^$ be tracially bounded maps with $\sup_{i \in I} \|u_i\|_{tb} \leq 1$. Let $\mathcal{U}$ be an ultrafilter on I and let $\hat{E}$ (resp. $\hat{F}$) be the ultraproduct of $(E_i)_{i \in I}$ (resp. $(F_i)_{i \in I}$) modulo $\mathcal{U}$. Let $\hat{u}: \hat{E} \rightarrow (\hat{F})^*$ be the map associated to the family $(u_i)_{i \in I}$. Then for all finite sets (x_k) in $\hat{E}$ and (y_k) in $\hat{F}$ we have*

$$(1.15) \quad \left| \sum \langle \hat{u}(x_k), y_k \rangle \right| \leq C \left\| \sum \lambda(g_k) \otimes x_k \right\|_{\min} \left\| \sum \lambda(g_k) \otimes y_k \right\|_{\min}.$$

Proof. We use the fact that $C_\lambda^*(F_\infty)$ is an exact C^* -algebra since it has the slice map property (*cf.* [DCH, Corollary 3.12] and see [Kr] for more details). Therefore if $F = \text{span}(\lambda(g_1), \dots, \lambda(g_n))$ we have $d_{SK}(F) = 1$. Then it is easy to derive (1.15) from (1.2) (applied to each u_i) taking (1.14) into account. ■

§2. The non-separability of OS_n .

We will denote by OS_n the set of all n dimensional operator spaces. We identify two elements $E, F \in OS_n$ if they are completely isometric. For $E, F \in OS_n$, let

$$d_{cb}(E, F) = \inf\{\|u\|_{cb}\|u^{-1}\|_{cb} \mid u: E \rightarrow F, u \text{ complete isomorphism}\}.$$

Then it can be shown (see [P1] for the easy details) that

$$\delta_{cb}(E, F) = \text{Log } d_{cb}(E, F)$$

is a distance on OS_n for which it is a complete metric space.

We will need a weaker metric structure on the space OS_n . To introduce it we need the following notation: For any linear map $u: E \rightarrow F$ between operator spaces we denote

$$\|u\|_k = \|I_{M_k} \otimes u\|_{M_k(E) \rightarrow M_k(F)}.$$

Note that

$$\|u\|_{cb} = \sup_{k \geq 1} \|u\|_k.$$

Now consider $E, F \in OS_n$. We define

$$d_k(E, F) = \inf\{\|u\|_k\|u^{-1}\|_k \mid u: E \rightarrow F, u \text{ linear isomorphism}\}.$$

Then by a simple compactness argument (the unit ball of the space $\mathcal{L}(E, F)$ of all linear maps is compact for any norm on $\mathcal{L}(E, F)$) one can check that

$$(2.1) \quad d_{cb}(E, F) = \sup_{k \geq 1} d_k(E, F).$$

We set

$$\delta_w(E, F) = \sum_{k \geq 1} 2^{-k} \text{Log } d_k(E, F).$$

Then, δ_w is a distance on OS_n . Let $\{E_i\}$ be a sequence in OS_n . Then $\delta_w(E_i, E) \rightarrow 0$ iff

$$d_k(E_i, F) \rightarrow 1 \quad \text{for all } k \geq 1.$$

In that case we will write simply $E_i \xrightarrow{w} E$. It was observed in [P1] that $E_i \xrightarrow{w} E$ iff for any non-trivial ultrafilter $\mathcal{U}$ on $\mathbf{N}$ the ultraproduct $\prod E_i / \mathcal{U}$ is completely isometric to E .

We will need the following known fact.

Proposition 2.1. *For any E, F in OS_n and any $k \geq 1$ we have*

$$d_{cb}(E, F) = d_{cb}(E^*, F^*)$$

and

$$d_k(E, F) = d_k(E^*, F^*)$$

for all $k \geq 1$. Hence in particular

$$\delta_{cb}(E, F) = \delta_{cb}(E^*, F^*) \quad \text{and} \quad \delta_w(E, F) = \delta_w(E^*, F^*).$$

Proof. It clearly suffices to know that for any $u: E \rightarrow F$ we have

$$(2.2) \quad \|u\|_{cb} = \|u^*\|_{cb}$$

and

$$(2.3) \quad \forall k \geq 1 \quad \|u\|_k = \|u^*\|_k.$$

The identity (2.2) was proved in [BP, ER1], while (2.3) is easy to check using the definition of $\|u\|_k$. We have (by [Sm])

$$(2.4) \quad \|u\|_k = \sup\{\|bua\|_{cb} \mid a: M_k^* \rightarrow E, b: F \rightarrow M_k, \|a\|_{cb} \leq 1, \|b\|_{cb} \leq 1\}.$$

Clearly (2.4) implies (2.3). Then the above proposition is obvious. ■

The following was proved in [P1].

Proposition 2.2. *Let $E \in OS_n$. The following are equivalent*

- (i) *For any sequence $\{E_i\}$ in OS_n tending weakly to E we have $d_{cb}(E, E_i) \rightarrow 1$ when $i \rightarrow \infty$.*
- (ii) *Same as (i) with each E_i isometric to E .*
- (iii) $d_{SK}(E) = d_{SK}(E^*) = 1$.

Remark. For any fixed integer $k \geq 1$, these are also equivalent to the same property as (i) restricted to E_i k -isometric to E .

Proof. We only prove (i) $\Rightarrow$ (iii) which is what we use in the sequel.

Assume (i). Then $E \subset B(\ell_2)$. Let $P_m: B(\ell_2) \rightarrow M_m$ be the projection which maps e_{ij} to itself if $1 \leq i, j \leq m$ and to zero otherwise. Let $E_m = P_m(E) \subset M_m$. It is very easy to check that $E_m \xrightarrow{w} E$. Hence if (i) holds we have

$$\begin{aligned} d_{SK}(E) &\leq d_{cb}(E, E_m) d_{SK}(E_m) \\ &\leq d_{cb}(E, E_m) \end{aligned}$$

$$\text{hence} \quad d_{SK}(E) \leq \lim d_{cb}(E, E_m) = 1.$$

Now if we apply the same to E^* (equipped with the dual operator space structure) we obtain by Proposition 2.1 that $d_{SK}(E^*) = 1$.

Reformulated in more concise terms, the proof reduces to this: let $OS_n(m)$ be the subset of OS_n formed of all n -dimensional subspaces of M_m . Then the union $\bigcup_{m \geq n} OS_n(m)$ is weakly dense in OS_n . Hence if (i) holds, E (and also E^* by Proposition 2.1) must be in the strong closure of $\bigcup_{m \geq n} OS_n(m)$, which means that $d_{SK}(E) = 1$ (and $d_{SK}(E^*) = 1$).

We can now prove

Theorem 2.3. *The metric space (OS_n, δ_{cb}) is non-separable if $n > 2$.*

Proof. Let $f: (OS_n, \delta_w) \longrightarrow (OS_n, \delta_{cb})$ be the identity mapping. Note that f^{-1} is continuous, but in general f is not.

However, if we assume OS_n strongly separable then we claim that f is in the first Baire class. Indeed, by (2.1) for any closed ball β in (OS_n, δ_{cb}) , $f^{-1}(\beta)$ is weakly closed. Hence if OS_n is strongly separable, for any U strongly open in OS_n $f^{-1}(U)$ must be an F_σ -set in weak topology, hence f is in the first Baire class. Note that the domain of f is compact hence is a Baire space. By Baire's classical theorem (cf. [Ba], see also [Ku, 31, X, Th. 1, p. 394]), if the range of f is separable, the set of points of continuity of f must be dense in the domain of f , i.e. dense for the weak topology.

This implies by Proposition 2.2 that for any E in OS_n there is a sequence $\{E_i\}$ in OS_n such that $d_{SK}(E_i) = d_{SK}(E_i^*) = 1$ which tends weakly to E . Equivalently, E can be viewed as the ultraproduct of (E_i) with respect to a non-trivial ultrafilter $\mathcal{U}$.

By Corollary 1.10 applied to the identity of E (with $u_i = I_{E_i}$), this implies that for any biorthogonal system $(x_1, \dots, x_n)$ $(x_1^*, \dots, x_n^*)$ in E we have

$$(2.5) \quad n \leq \left\| \sum_1^n \lambda(g_i) \otimes x_i \right\|_{C_\lambda \otimes_{\min} E} \left\| \sum \lambda(g_i) \otimes x_i^* \right\|_{C_\lambda \otimes_{\min} E^*}.$$

Now let $E_n^\lambda = \text{span}(\lambda(g_1), \dots, \lambda(g_n)) \subset C_\lambda$.

Let $\lambda_*(g_i)$ be the biorthogonal functionals in $(E_n^\lambda)^*$. Then, if $n > 1$, we have by [AO]

$$\left\| \sum_1^n \lambda(g_i) \otimes \lambda(g_i) \right\| = 2\sqrt{n-1},$$

and since $t = \sum_{i=1}^n \lambda(g_i) \otimes \lambda_*(g_i)$ represents the inclusion map $j: E_n^\lambda \rightarrow C_\lambda$, we have

$$\|t\|_{C_\lambda \otimes_{\min} (E_n^\lambda)^*} = \|j\|_{cb(E_n^\lambda, C_\lambda)} = 1.$$

Hence taking $x_i = \lambda(g_i)$, $x_i^* = \lambda_*(g_i)$ in (2.5) we obtain

$$n \leq 2\sqrt{n-1}$$

or equivalently $n \leq 2$. ■

Remark 2.4. By a simple modification of the preceding proof, one can prove that the subset $HOS_n \subset OS_n$ formed of all the n -dimensional operator spaces which are isometric to ℓ_2^n is non-separable if $n > 2$. Our original argument here gave only $n > 4$, the improvement is due to Timur Oikhberg.

Here is briefly the argument: By the proof of Theorem 2.3, if HOS_n is separable, then any E in HOS_n must satisfy (2.5). Consider then the operator space $\min(\ell_2^n)$ obtained by embedding ℓ_2^n isometrically into a commutative C^* -algebra. Let (e_i) be the basis of ℓ_2^n .

Assume $n > 1$. Let $\gamma_n = 2(1 - n^{-1})^{1/2}$. Consider the subspace

$$E \subset E_n^\lambda \oplus \min(\ell_2^n)$$

spanned by the vectors $x_i = \lambda(g_i) \oplus \gamma_n e_i$. Then by [AO, p. 1038] we have $\|\sum \alpha_i x_i\| = \gamma_n (\sum |\alpha_i|^2)^{1/2}$,

$\forall (\alpha_i) \in \mathbb{C}^n$, hence $E \in HOS_n$. Furthermore, we have

$$(2.6) \quad \left\| \sum_1^n \lambda(g_i) \otimes x_i \right\|_{C_\lambda \otimes_{\min} E} = \max\{2\sqrt{n-1}, \gamma_n^2\} \quad \text{and} \quad \left\| \sum \lambda(g_i) \otimes x_i^* \right\|_{C_\lambda \otimes_{\min} E^*} \leq 1,$$

so that we conclude again from (2.5) that $n \leq 2$.

Remark 2.5. (i) An operator space E is called homogeneous if for any $u: E \rightarrow E$ we have $\|u\| = \|u\|_{cb}$. This notion seems particularly interesting in the Hilbertian case (see [P3]). Consider an arbitrary n -dimensional operator space E given with a basis $(e_1, \dots, e_n)$. We can define its "homogeneous hull" $\hat{E}$ as follows. Let $U(n)$ be the unitary group. We view the coordinates (u_{ij}) of a unitary matrix u as a continuous function on $U(n)$, so that $u_{ij} \in C(U(n))$. In the space $C(U(n)) \otimes_{\min} E$ we consider the elements

$$\hat{e}_i = \sum_j u_{ij} \otimes e_j \in C(U(n)) \otimes_{\min} E.$$

Let $\hat{E}$ be the operator space spanned by $(\hat{e}_1, \dots, \hat{e}_n)$. Observe that $U(n)$ acts isometrically (and actually completely isometrically) on $\hat{E}$, therefore it is easy to check that $\hat{E}$ is Hilbertian and homogeneous. Moreover, we have $d_{SK}(\hat{E}) \leq d_{SK}(E)$. Now let F be another operator space and let $u: E \rightarrow F$ be an isomorphism. Let $f_i = u(e_i)$ and let $\hat{F}$ be the operator space associated to F and this basis. Then it is easy to check that

$$d_{cb}(\hat{E}, \hat{F}) \leq \|u\|_{cb} \|u^{-1}\|_{cb} \quad \text{and} \quad d_k(\hat{E}, \hat{F}) \leq \|u\|_k \|u^{-1}\|_k \quad \forall k.$$

(ii) Let us denote by HH_n the subset of OS_n formed of all the Hilbertian homogeneous spaces. Using the first part of this remark, it is easy to check that any space E in HH_n is the weak limit of a net (E_i) in HH_n such that $d_{SK}(E_i) = 1$ for all i . Then a simple modification of the proof of Theorem 2.3 shows that HH_n is a non-separable subset of OS_n for the (strong) distance δ_{cb} , if $n > 2$. Indeed, we can replace the space E and its basis (x_i) in Remark 2.4 by $\hat{E}$ and $\hat{x}_i$. Using an inequality due to Haagerup [H3, Lemma 2.4], one can check that (2.6) remains valid. This gives us a space in HH_n which satisfies (2.5) only if $n \leq 2$. (Alternately, one could replace E by the linear span of a circular system in the sense of Voiculescu [VDN], but this seems to yield non-separability only for $n > 4$.)

Proposition 2.6. *Let A be any separable C^* -algebra (or any separable operator space) and let $n \geq 1$.*

- (i) *The subset $S_n(A) \subset OS_n$ formed of all the n -dimensional subspaces of A is separable in (OS_n, δ_{cb}) .*

- (ii) For any $n > 2$, there is an operator space $E_0 \in OS_n$ and $\varepsilon_0 > 0$ such that for any n -dimensional subspace $E \subset A$ we have $d_{cb}(E, E_0) \geq 1 + \varepsilon_0$.

Proof. The first part is proved by a standard perturbation argument. We merely sketch it: Let $D \subset A$ be a dense countable subset. For any n -tuple $x = (x_1, \dots, x_n)$ of linearly independent elements of A , let $E_x \subset A$ be their linear span. Let $x_1^*, \dots, x_n^*$ be functionals in A^* which are biorthogonal to x_i . Fix $\varepsilon > 0$. Pick $y_1, \dots, y_n$ in D such that $\sum \|x_i - y_i\| < \varepsilon$. Consider then the operator $u: A \rightarrow A$ associated to $\sum x_i^* \otimes (y_i - x_i)$. Clearly $\|u\|_{cb} \leq f(\varepsilon)$ with $f(\varepsilon) \rightarrow 0$ when $\varepsilon \rightarrow 0$. Moreover, $(I + u)(x_i) = y_i$, and (say, when $f(\varepsilon) < 1$) $x_i = (I + u)^{-1}(y_i)$. This immediately yields $d_{cb}(E_x, E_y) \leq 1 + g(\varepsilon)$ with $g(\varepsilon) \rightarrow 0$ when $\varepsilon \rightarrow 0$, whence (i).

The second part follows from the first one, since it merely expresses the fact that (by Theorem 2.3) $S_n(A)$ is not dense in OS_n for $n > 2$. ■

We now give a more precise version of Theorem 2.3 based on the following well known variant of Baire's Theorem.

Lemma 2.7. *Let S, T be metric spaces and let $f: S \rightarrow T$ be a mapping such that for every closed ball $B \subset T$, $f^{-1}(B)$ is closed in S . Fix a number $\varepsilon > 0$. Let $C_\varepsilon(f)$ be the set of all points of ε -continuity of f , i.e. all points s in S such that whenever $s_i \rightarrow s$ we have $\limsup d(f(s_i), f(s)) \leq \varepsilon$. Now assume that T is ε -separable, i.e. there exists a sequence $\{B_n\}$ of closed balls of radius ε in T such that $T = \cup B_n$. Then, if S is a Baire space, the set $C_{2\varepsilon}(f)$ of points of (2ε) -continuity of f is dense in S .*

Proof. Let $S' = \bigcup_n [f^{-1}(B_n) \setminus \overset{\circ}{f^{-1}(B_n)}] \subset S$. Since S' is a countable union of closed sets with empty interior in a Baire space, its complement $S \setminus S'$ is dense in S . But it is easy to check (recall $S = \cup f^{-1}(B_n)$), that $S \setminus S' \subset C_{2\varepsilon}(f)$. ■

Theorem 2.8. *For each $n \geq 1$, let ε_n be the infimum of all numbers $\varepsilon > 0$ such that (OS_n, δ_{cb}) is ε -separable. Let $\delta_n = \exp(\varepsilon_n)$. Then we have for all $n \geq 3$*

$$(2.7) \quad \left(\frac{n}{2\sqrt{n-1}} \right)^{1/4} \leq \delta_n.$$

and there is a constant $c > 0$ such that for all $n \geq 3$

$$(2.8) \quad cn^{1/8} < \delta_n \leq n^{1/2}.$$

Proof. We apply Lemma 2.7 to the same map f as in the proof of Theorem 2.3.

Consider $\delta > \delta_n$ and let $\varepsilon = \log(\delta)$ so that (OS_n, δ_{cb}) is ε -separable. By Lemma 2.7, $C_{2\varepsilon}(f)$ is δ_w -dense. Then, by a simple modification of the proof of Proposition 2.2, any E is OS_n is the weak limit of a net E_i such that $\limsup d_{SK}(E_i) \leq e^{2\varepsilon} = \delta^2$ and $\limsup d_{SK}(E_i^*) \leq e^{2\varepsilon} = \delta^2$. Then (recalling Corollary 1.10) we conclude, as in the proof of Theorem 2.3, that

$$n \leq \delta^4 2\sqrt{n-1}$$

whence (2.7).

It was proved in [P3, Theorem 9.6] that $d_{SK}(E) \leq n^{1/2}$ for all E in OS_n , therefore $\delta_n \leq n^{1/2}$ and (2.8) follows. ■

Corollary 2.9. *There is a constant $c > 0$ such that for each $n \geq 3$ there is an uncountable collection $(E_i)_{i \in I}$ is OS_n satisfying*

$$\forall i \neq j \quad d_{cb}(E_i, E_j) > cn^{1/8}$$

It would be interesting to find the exact asymptotic behaviour of δ_n .

Remark 2.10. Two or three months after this paper had been circulated as a preprint, Simon Wassermann mentioned to us that he conjectured that the linear spans of the n -tuples of operators considered by Voiculescu in [V] should yield a more explicit non-separable family of finite dimensional operator spaces. Some form of this conjecture is indeed correct. This shows that groups with Kazhdan's property T (see [DHV]) can be used to prove the non-separability of OS_n . Here are the details. In [V], to each subset Ω of the integers, Voiculescu associates an n -tuple $\tau_\Omega = (T_1^\Omega, \dots, T_n^\Omega)$ of operators in $B(H)$ (with say $H = \ell_2$), in the following way. Let G be any discrete group with Kazhdan's property T admitting a countable collection of pairwise disjoint finite dimensional representations (π_k) . (For instance we can take $G = SL_3(\mathbb{Z})$.) Let $(g_1, \dots, g_{n-1})$ be a finite set of generators and let g_n be equal to the unit element. Then for any $\Omega \subset \mathbb{N}$, we define

$$T_j^\Omega = \oplus_{k \in \Omega} \pi_k(g_j), \quad j = 1, 2, \dots, n.$$

Let M_Ω be the finite dimensional operator space spanned by τ_Ω . We claim these form a non-separable collection of operator spaces. This does not seem to follow from Voiculescu's stated results but it does follow easily from the proof of his proposition 3. Indeed he shows a stronger result than he states, as follows:

First, in case M_Ω is not n -dimensional we consider an n -dimensional operator space E_Ω containing τ_Ω .

Following [V], we use property T through the following: There is a fixed number $\varepsilon > 0$ such that if Ω and Ω' are subsets of $\mathbb{N}$ with $\Omega \not\subset \Omega'$, then there are unitary operators $u_1, \dots, u_n$ satisfying

$$(2.9) \quad n = \left\| \sum_1^n T_j^\Omega \otimes u_j \right\|_{\min} \quad \text{and} \quad \left\| \sum_1^n T_j^{\Omega'} \otimes u_j \right\|_{\min} < n - \varepsilon.$$

(More precisely, if we pick $k \in \Omega - \Omega'$ then we can take $u_j = \overline{\pi_k(g_j)}$.)

Now assume that the metric space (OS_n, δ_{cb}) is separable. Let (E_m) be a dense sequence in OS_n . Fix a number $\eta > 0$. Then there is an integer m and a continuous collection $\mathcal{C}$ of subsets of $\mathbb{N}$ such that for each Ω in $\mathcal{C}$ there is a map $v_\Omega: E_\Omega \rightarrow E_m$ such that $\|v_\Omega\|_{cb} < 1 + \eta$ and $\|v_\Omega^{-1}\|_{cb} = 1$. Now consider the continuous family $(v_\Omega(T_j^\Omega))_{j \leq n}$ of n -tuples of elements of E_m . Since $(E_m)^n$ (the space of n -tuples of elements of E_m) is norm-separable, there must exist a continuous subcollection $\mathcal{C}_1 \subset \mathcal{C}$ such that for all Ω, Ω' in $\mathcal{C}_1$ we have

$$(2.10) \quad \sum_1^n \|v_\Omega(T_j^\Omega) - v_{\Omega'}(T_j^{\Omega'})\| < \eta.$$

A fortiori, $\mathcal{C}_1$ has cardinality > 1 , hence we can find Ω, Ω' in $\mathcal{C}_1$ satisfying $\Omega \not\subset \Omega'$. By (2.9) this implies

$$\left\| \sum v_{\Omega'}(T_j^{\Omega'}) \otimes u_j \right\|_{\min} \leq \|v_{\Omega'}\|_{cb} \left\| \sum T_j^{\Omega'} \otimes u_j \right\|_{\min} \leq (1 + \eta)(n - \varepsilon),$$

and

$$n = (\|v_\Omega^{-1}\|_{cb})^{-1} \left\| \sum_1^n T_j^\Omega \otimes u_j \right\|_{\min} \leq \left\| \sum v_\Omega(T_j^\Omega) \otimes u_j \right\|_{\min}.$$

This gives by (2.10)

$$n \leq \left\| \sum v_{\Omega'}(T_j^{\Omega'}) \otimes u_j \right\|_{\min} + \sum_1^n \|v_{\Omega'}(T_j^{\Omega'}) - v_\Omega(T_j^\Omega)\| \leq (1 + \eta)(n - \varepsilon) + \eta.$$

When $\eta > 0$ is small enough this is impossible (since $\varepsilon > 0$ remains fixed). This contradiction completes the proof. ■

Actually, Simon Wassermann conjectured that the whole family of spaces (M_Ω) is uniformly d_{cb} -separated, i. e. that for some $\eta > 0$ we have $d_{cb}(M_\Omega, M_{\Omega'}) > 1 + \eta$ whenever $\Omega \neq \Omega'$. As far as we know this is still open.

It is known ([Tr]) that $G = SL_3(\mathbb{Z})$ admits two generators so that (recall that g_n is the unit) we obtain by this reasoning a continuous collection of 3-dimensional operator spaces (E_t) such that for some $\varepsilon > 0$ we have $d_{cb}(E_t, E_s) > 1 + \varepsilon$ for all $t \neq s$. By a simple modification, we can make sure that the spaces we obtain are spanned by three unitaries. The same cannot be achieved with spans of two unitaries. Indeed, the span of two unitaries (U_1, U_2) is completely isometric to the span of (I, U) with $U = U_1^* U_2$, which itself embeds completely isometrically into a commutative (hence nuclear) C^* -algebra.

However, this approach does not seem to yield the refinements in Remarks 2.4 and 2.5 or in Corollary 2.9.

§3. Applications to $B(H) \otimes B(H)$ and maximal operator spaces.

By Kirchberg's results in [Ki2], Theorem 2.3 implies

Corollary 3.1. *If $\dim H = \infty$, there is more than one C^* -norm on $B(H) \otimes B(H)$. In other words we have*

$$B(H) \otimes_{\min} B(H) \neq B(H) \otimes_{\max} B(H).$$

Proof. This follows from the equivalence of the conjectures (A7) and (A2) in [Ki2, p. 483]. For the convenience of the reader, we include a direct argument, as follows. For any discrete group G we denote by $C^*(G)$ the full C^* -algebra of G . By Proposition 2.6 (ii) (applied with $A = C^*(F_\infty)$), there is an operator space E_0 such that for some $\varepsilon_0 > 0$ we have

$$d_{cb}(E, E_0) \geq 1 + \varepsilon_0$$

for all n -dimensional subspaces $E \subset C^*(F_\infty)$. Now let F_I be a free group associated to a set of generators $\{g_i \mid i \in I\}$ where I is any set with infinite cardinality. Observe that for any finite dimensional (or merely separable) subspace $E \subset C^*(F)$ there is a countable infinite subset $J \subset I$ such that $E \subset C^*(F_J)$. (Indeed only countably many “letters” are being used.) Hence we also have

$$d(E, E_0) \geq 1 + \varepsilon_0$$

for all n -dimensional subspaces $E \subset C^*(F_I)$. Let $\pi: C^*(F) \rightarrow B(H)$ be a C^* -algebra representation of the full C^* -algebra of a big enough free group F onto $B(H)$. We have

$$B(H) \approx C^*(F)/\text{Ker } \pi.$$

Let $\mathcal{I} = \text{Ker } \pi$. Then (by Lemma 0.8) the quotient norm of the space

$$Q = [C^*(F) \otimes_{\min} B(H)] / [\mathcal{I} \otimes_{\min} B(H)]$$

induces on $B(H) \otimes B(H)$ a C^* -norm. Assume that there is only one such norm. Then we have isometrically

$$(3.1) \quad Q = B(H) \otimes_{\min} B(H).$$

Now we may clearly assume that E_0^* is a subspace of $B(H)$, i.e. we have $E_0^* \subset B(H)$ completely isometrically, so that the completely isometric inclusion $j: E_0 \rightarrow B(H)$ can be viewed as an element j_0 in $B(H) \otimes_{\min} E_0^* \subset B(H) \otimes_{\min} B(H)$. Note that $\|j_0\|_{\min} = \|j\|_{cb} = 1$. By Lemma 0.7 and by (3.1), for any $\varepsilon > 0$ there is a lifting $\tilde{j}_0$ in $C^*(F) \otimes_{\min} E_0^*$ with $\|\tilde{j}_0\|_{\min} < 1 + \varepsilon$ and $(\pi \otimes I_{E_0^*})(\tilde{j}_0) = j_0$. Now, let $\tilde{j}: E_0 \rightarrow C^*(F)$ be the associated linear operator. We have $\pi\tilde{j} = j$ and $\|\tilde{j}_0\|_{\min} = \|\tilde{j}\|_{cb}$, hence $d_{cb}(E_0, \tilde{j}(E_0)) \leq \|\pi\|_{cb}\|\tilde{j}\|_{cb} < 1 + \varepsilon$. When $\varepsilon < \varepsilon_0$ this is impossible. $\blacksquare$

Remark. By [Ki2, section 8], Corollary 3.1 has the following consequences

(i) There is a separable unital C^* -algebra with the WEP in the sense of Lance [La] for which $\text{Ext}(A)$ is not a group.

(ii) There are separable unital C^* -algebras A, B with WEP such that

$$A \otimes_{\min} B \neq A \otimes_{\max} B.$$

(iii) The WEP does not imply the local lifting property (in short LLP) in the sense of [Ki2].

(iv) There is a separable unital C^* -algebra A with WEP which is not approximately injective in the sense of [EH].

(v) The identity $A^{op} \otimes_{\min} A = A^{op} \otimes_{\max} A$ does not imply the approximate injectivity of A .

(vi) There is a unital separable C^* -algebra B with WEP which is not a quotient C^* -algebra of an approximately injective C^* -algebra.

By [Ki2, p. 484], Corollary 3.1 also implies a negative answer to Kirchberg's conjecture (C1) in [Ki2]. Thus we have :

$C^*(F_\infty)$ and $C^*(SL_2(\mathbb{Z}))$ are not approximately injective in $B(H)$.

Finally, by [Ki2, p. 487], Corollary 3.1 implies a negative answer to the conjecture (P2) in [Ki2], hence we have : Let $K = K(H)$, $B = B(H)$. Consider the canonical morphism

$$\Phi: B \otimes_{\min} B \mapsto (B/K) \otimes_{\min} (B/K).$$

Then the kernel of Φ is strictly larger than the set $F(K, B, B \otimes_{\min} B) + F(B, K, B \otimes_{\min} B)$ where $F(., ., .)$ denotes the Fubini product.

In a different direction, we can give more examples of non-exact operator spaces, completing those of [P1]. Following Blecher and Paulsen [BP], given a Banach space E , we denote by $\min(E)$ the operator space obtained by embedding E into a commutative C^* -algebra, or equivalently by embedding E into the space $C(K)$ of all continuous functions on K with $K = (B_{E^*}, \sigma(E^*, E))$. Similarly, let I be a suitable set and let $g: \ell_1(I) \rightarrow E$ be a metric surjection (i.e. q^* is an isometry). We view the space $\ell_1(I) = c_0(I)^*$ as an operator space with the dual operator space structure. Then (cf. [BP, Pa2]) we denote by $\max(E)$ the operator space obtained by equipping E with the operator space structure (in short o.s.s.) of the quotient space $\ell_1(I)/\text{Ker}(g)$. Equivalently, we have a complete isometry

$$\max(E) \longrightarrow (\min(E^*))^*,$$

and $\max(E)$ is characterized by the isometric identity

$$(3.2) \quad cb(\max(E), M_n) = B(E, M_n).$$

More generally, for any operator space F we have isometrically

$$(3.2)' \quad cb(\max(E), F) = B(E, F).$$

We refer to [BP, Pa2] for more information. It will be convenient to introduce the following characteristic for an operator space E .

$$d_{QSK}(E) = \inf\{d_{cb}(E, F)\}$$

where the infimum runs over all operator spaces F of the form $F = E_1/E_2$ where $E_2 \subset E_1 \subset K$ and $\dim E = \dim F$.

Theorem 3.2. *Let E be any n -dimensional Banach space. Then*

$$d_{QSK}(\max(E)) \geq \frac{\sqrt{n}}{4}.$$

Proof. By a well known result in Banach space theory (cf. e.g. [P2, Theorem 1.11 p.15]) we have $\pi_2(I_{E^*}) = \sqrt{n}$. Let $u: X \rightarrow Y$ be a 2-absolutely summing operator between Banach spaces and let $J: Y \rightarrow Y_1$ be an isometric embedding. Then it is easy to see that

$$(3.3) \quad \pi_2(u) = \pi_2(Ju).$$

Consider an isomorphism $v: E_1/E_2 \rightarrow \max(E)$ with $E_2 \subset E_1 \subset K$. Observe $(E_1/E_2)^* = E_2^\perp \subset E_1^*$. Let $J: E_2^\perp \rightarrow E_1^*$ be the (isometric) canonical inclusion. We will now apply Corollary 1.7 to the map

$$Jv^*: \min(E^*) \xrightarrow{v^*} E_2^\perp \xrightarrow{J} E_1^*.$$

Note that $d_{SK}(\min(E^*)) = d_{SK}(E_1) = 1$. Hence Corollary 1.7 yields by (3.3)

$$\pi_2(v^*) = \pi_2(Jv^*) \leq 4\|Jv^*\|_{cb} \leq 4\|v^*\|_{cb} = 4\|v\|_{cb}$$

but this implies by (0.4)

$$n^{1/2} = \pi_2(I_{E^*}) \leq \|v^{-1*}\| \pi_2(v^*) \leq 4\|v^{-1}\| \|v\|_{cb}$$

hence $n^{1/2} \leq 4d_{QSK}(\max(E))$. ■

Remark. In particular the preceding result answers a question raised by Vern Paulsen (private communication): the space $\max(\ell_2^n)$ is quite different (when n is large enough) from the linear span of n Clifford matrices, i.e. matrices (u_i) in M_{2^n} satisfying the relations

$$u_i = u_i^* \quad u_i u_j^* + u_j^* u_i = 2\delta_{ij} I.$$

Indeed, if we denote $Cl_n = \text{span}(u_1, \dots, u_n)$ then by Theorem 3.2 we have

$$d_{cb}(\max(\ell_2^n), Cl_n) \geq d_{QSK}(\max(\ell_2^n)) \geq \sqrt{n}/4.$$

§4. A new tensor product for C^* -algebras or operator spaces.

Let $E_1 \subset B(H_1), E_2 \subset B(H_2)$ be arbitrary operator spaces.

Let us denote by $\|\cdot\|_M$ the norm induced on the algebraic tensor product $E_1 \otimes E_2$ by $B(H_1) \otimes_{\max} B(H_2)$. Recalling (0.3), and using the injectivity of $B(H_1)$ and $B(H_2)$ as well as the decomposition property of *c.b.* maps into $B(H)$, it is easy to check (see Lemma 4.1 below) that this norm is independent of the choice of the completely isometric embeddings $E_1 \subset B(H_1), E_2 \subset B(H_2)$. In other words, the norm $\|\cdot\|_M$ on $E_1 \otimes E_2$ depends only on the operator space structures of E_1 and E_2 .

We will denote by $E_1 \otimes_M E_2$ the completion of $E_1 \otimes E_2$ under this norm. We equip the space $E_1 \otimes_M E_2$ with the natural operator space structure induced by the C^* -algebra $B(H_1) \otimes_{\max} B(H_2)$ via the isometric embedding $E_1 \otimes_M E_2 \subset B(H_1) \otimes_{\max} B(H_2)$.

Clearly, if A, B are C^* -algebras, then $\|\cdot\|_M$ is a C^* -norm on $A \otimes B$ and $A \otimes_M B$ also is a C^* -algebra.

Lemma 4.1. *Let F_1, F_2 be two operator spaces. Consider *c.b.* maps $u_1 : E_1 \rightarrow F_1$ and $u_2 : E_2 \rightarrow F_2$. Then $u_1 \otimes u_2$ defines a *c.b.* map from $E_1 \otimes_M E_2$ to $F_1 \otimes_M F_2$ with*

$$(4.1) \quad \|u_1 \otimes u_2\|_{cb(E_1 \otimes_M E_2, F_1 \otimes_M F_2)} \leq \|u_1\|_{cb} \|u_2\|_{cb}.$$

Proof. Indeed, note that if $F_1 \subset B(\mathcal{H}_1), F_2 \subset B(\mathcal{H}_2)$ then by the extension property of *c.b.* maps (*cf.* [Pa1] p. 100) u_1, u_2 admit extensions $\tilde{u}_1 : B(H_1) \rightarrow B(\mathcal{H}_1)$ and $\tilde{u}_2 : B(H_2) \rightarrow B(\mathcal{H}_2)$ with $\|\tilde{u}_1\|_{cb} = \|u_1\|_{cb}$ and $\|\tilde{u}_2\|_{cb} = \|u_2\|_{cb}$. Hence it suffices to check this in the case when each of E_1, E_2, F_1, F_2 is $B(H)$ for some H . Then the idea is to use the decomposition property of *c.b.* maps on $B(H)$ as linear combinations of completely positive maps to reduce checking (4.1) to the case of completely positive maps. In the completely positive case, the relevant point here is of course (0.3). This idea leads to a simple proof of (4.1) with some additional numerical factor. However, this factor can be removed at the cost of a slightly more technical argument based on [H3]. For lack of a suitable reference, we now briefly outline this (straightforward) argument to check (4.1). Let A, B be C^* -algebras, we will denote by $CP(A, B)$ the set of all completely positive maps $u : A \rightarrow B$, and by $D(A, B)$ the set of all decomposable maps $u : A \rightarrow B$, *i.e.* maps which can be written as $u = u_1 - u_2 + i(u_3 - u_4)$ with $u_1, \dots, u_4 \in CP(A, B)$.

In [H3], Haagerup defines the norm $\|u\|_{dec}$ on $D(A, B)$ as follows. Consider all possible mappings S_1, S_2 in $CP(A, B)$ such that the map $v: A \rightarrow M_2(B)$ defined by

$$v(x) = \begin{pmatrix} S_1(x) & u(x^*)^* \\ u(x) & S_2(x) \end{pmatrix}$$

is completely positive. Then we set $\|u\|_{dec} = \inf\{\max\{\|S_1\|, \|S_2\|\}\}$ where the infimum runs over all possible such mappings.

In [H3, Proposition 1.3 and Theorem 1.6] the following results appear:

$$(4.2) \quad \forall u \in D(A, B) \quad \|u\|_{cb} \leq \|u\|_{dec},$$

$$(4.3) \quad \forall u \in CP(A, B) \quad \|u\| = \|u\|_{cb} = \|u\|_{dec},$$

(see also [Pa1, p. 28] for the first equality)

(4.4) if C is any C^* -algebra, if $u \in D(A, B)$ and $v \in D(B, C)$, then $vu \in D(A, C)$ and

$$\|vu\|_{dec} \leq \|v\|_{dec}\|u\|_{dec},$$

$$(4.5) \quad \forall u \in cb(B(H), B(\mathcal{H})) \quad \|u\|_{cb} = \|u\|_{dec}.$$

Now let A_1, A_2, B_1, B_2 be arbitrary C^* -algebras, and let $u_1 \in D(A_1, B_1)$, $u_2 \in D(A_2, B_2)$. We claim that Haagerup's results imply that $u_1 \otimes u_2$ extends to a decomposable map from $A_1 \otimes_{\max} A_2$ into $B_1 \otimes_{\max} B_2$ (still denoted by $u_1 \otimes u_2$) satisfying

$$(4.6) \quad \|u_1 \otimes u_2\|_{dec} \leq \|u_1\|_{dec}\|u_2\|_{dec}.$$

By (4.2) we have a fortiori

$$(4.7) \quad \|u_1 \otimes u_2\|_{cb(A_1 \otimes_{\max} A_2, B_1 \otimes_{\max} B_2)} \leq \|u_1\|_{dec}\|u_2\|_{dec}.$$

To verify (4.6) (hence also (4.7)) we may assume (using (4.4) and $u_1 \otimes u_2 = (u_1 \otimes I)(I \otimes u_2)$) that $A_2 = B_2$ and u_2 is the identity on A_2 . Consider then $v_1: A_1 \rightarrow M_2(B_1)$ of the form

$$v_1(x) = \begin{pmatrix} S_1(x) & u_1(x^*)^* \\ u_1(x) & S_2(x) \end{pmatrix}$$

with v_1 and S_1, S_2 all completely positive.

By (0.3), the associated map $v_1 \otimes I_{A_2}$ as well as $S_1 \otimes I_{A_2}$ and $S_2 \otimes I_{A_2}$ are (bounded and) completely positive from $A_1 \otimes_{\max} A_2$ to $B_1 \otimes_{\max} A_2$. Therefore (by the definition of $\|\cdot\|_{dec}$)

$$\|u_1 \otimes I_{A_2}\|_{dec} \leq \max\{\|S_1 \otimes I_{A_2}\|, \|S_2 \otimes I_{A_2}\|\}$$

$$\text{hence by (0.3)} \quad \leq \max\{\|S_1\|, \|S_2\|\}.$$

It follows that

$$\|u_1 \otimes I_{A_2}\|_{dec} \leq \|u_1\|_{dec},$$

and this is enough to verify (4.6) (and a fortiori (4.7)).

The proof of (4.1) is now easy: let $u_1, \tilde{u}_1$ and $u_2, \tilde{u}_2$ be as explained above, by (4.7)

$$\begin{aligned} \|u_1 \otimes u_2\|_{cb(E_1 \otimes_M E_2, F_1 \otimes_M F_2)} &\leq \|\tilde{u}_1 \otimes \tilde{u}_2\|_{cb(B(H_1) \otimes_{\max} B(H_2), B(\mathcal{H}_1) \otimes_{\max} B(\mathcal{H}_2))} \\ &\leq \|\tilde{u}_1\|_{dec} \|\tilde{u}_2\|_{dec} \end{aligned}$$

$$\text{hence by (4.5)} \quad \leq \|\tilde{u}_1\|_{cb} \|\tilde{u}_2\|_{cb} = \|u_1\|_{cb} \|u_2\|_{cb}.$$

This completes the proof of (4.1). ■

We will use several times the following obvious consequence of (4.1) and the definition of $E_1 \otimes_M E_2$:

(4.8) If u_1 and u_2 (as above) are complete isometries, then $u_1 \otimes u_2: E_1 \otimes_M E_2 \rightarrow F_1 \otimes_M F_2$ also is a complete isometry.

By Corollary 3.1, we know that there are operator spaces E, F such that $E \otimes_{\min} F \neq E \otimes_M F$. It is natural to try to understand the meaning of this new tensor norm $\|\cdot\|_M$ and to characterize the operator spaces E, F for which the equality holds. For that purpose the following result due to Kirchberg [Ki3] will be crucial:

For any free group F_I and any H , we have an isometric identity

$$(4.9) \quad C^*(F_I) \otimes_{\min} B(H) = C^*(F_I) \otimes_{\max} B(H).$$

Using this, we have

Proposition 4.2. *Let E, F be operator spaces, let $u \in E \otimes F$ and let $U: F^* \rightarrow E$ be the associated finite rank linear operator. Consider a finite dimensional subspace $S \subset C^*(F_\infty)$ and a factorization of U of the form $U = ba$ with bounded linear maps $a: F^* \rightarrow S$ and $b: S \rightarrow E$, where $a: F^* \rightarrow S$ is weak-* continuous. Then*

$$\|u\|_M = \inf\{\|a\|_{cb}\|b\|_{cb}\}$$

where the infimum runs over all such factorizations of U .

Proof. Assume $E \subset B(H)$ and $F \subset B(K)$ with H, K Hilbert. It clearly suffices to prove this in the case when E and F are both finite dimensional. Assume U factorized as above with $\|a\|_{cb}\|b\|_{cb} < 1$. Then by Kirchberg's theorem (4.9) the min and max norms are equal on $C^*(F_\infty) \otimes B(K)$, hence, by (4.8), we have isometrically $S \otimes_{\min} F = S \otimes_M F$, so that if $\hat{a}$ is the element of $S \otimes_{\min} F$ associated to a , we have $\|\hat{a}\|_M = \|a\|_{cb}$ and $u = (b \otimes I_F)(\hat{a})$. Therefore, by (4.1) we have $\|u\|_M \leq \|b\|_{cb}\|\hat{a}\|_M \leq \|a\|_{cb}\|b\|_{cb} < 1$.

The proof of the converse is essentially the same as for Corollary 3.1 above. We skip the details. ■

Let A be any C^* -algebra. For any finite dimensional operator space E , let

$$d_{SA}(E) = \inf\{d_{cb}(E, F) \mid F \subset A\}.$$

Then the preceding result immediately implies

Corollary 4.3. *Let E be a finite dimensional operator space. Let $i_E \in E \otimes E^*$ be the tensor associated to the identity on E . Then*

$$\|i_E\|_M = d_{SC^*(F_\infty)}(E).$$

In particular we have

$$(4.10) \quad d_{SC^*(F_\infty)}(E) = d_{SC^*(F_\infty)}(E^*).$$

Proof. The first part is clear by Proposition 4.2. To check (4.10), observe more generally that for any operator spaces E, F , the "flip isomorphism" $(x \otimes y \rightarrow y \otimes x)$ is a complete isometry between the spaces $E \otimes_M F$ and $F \otimes_M E$. Hence (4.10) follows by symmetry. ■

Remark. The preceding argument shows the following : If $A \otimes_{\min} B(H) = A \otimes_{\max} B(H)$ then for any finite dimensional operator space E we have

$$(4.11) \quad d_{SC^*(F_\infty)}(E) \leq d_{SA}(E^*).$$

In particular, this holds (by definition of nuclearity) if A is nuclear. (Hence (4.11) still holds if A is exact, since $d_{SK}(E^*) \leq d_{SA}(E^*)$ in that case). The proof of Corollary 3.1 shows that (4.11) is false in general for $A = B(H)$. (Observe that $d_{SA}(E^*) = 1$ for any E if $A = B(H)$ and $\dim(H) = \infty$.)

Remark. We can now give a quantitative version of Corollary 3.1. For any n let

$$(4.12) \quad \lambda(n) = \sup \left\{ \frac{\|u\|_{\max}}{\|u\|_{\min}} \right\}$$

where the supremum runs over all u in $B(H) \otimes B(H)$ with $\text{rank}(u) \leq n$. We claim that (with the notation of Theorem 2.8)

$$(4.13) \quad (cn^{1/8} \leq) \quad \delta_n \leq \lambda(n) \leq \sqrt{n}.$$

To verify this, first observe that

$$(4.12)' \quad \lambda(n) = \sup \left\{ \frac{\|u\|_M}{\|u\|_{\min}} \right\}$$

where the supremum runs over all operator spaces E, F in OS_n and all u in $E \otimes F$. Equivalently, we have

$$(4.12)'' \quad \lambda(n) = \sup \{\|i_E\|_M\}$$

where the supremum runs over all E in OS_n and where $i_E \in E \otimes E^*$ represents (as above) the identity on E .

Indeed, any u as in (4.12)' can be rewritten as $u = (I_E \otimes \tilde{u})(i_E)$ where $\tilde{u}: E^* \rightarrow F$ is the linear map corresponding to u , hence by (4.1) we have $\|u\|_M \leq \|\tilde{u}\|_{cb}\|i_E\|_M = \|u\|_{\min}\|i_E\|_M$, and (4.12)'' follows.

By Corollary 4.3, this implies

$$(4.12)''' \quad \lambda(n) = \sup \{d_{SC^*(F_\infty)}(E) \mid E \in OS_n\}.$$

Now since $C^*(F_\infty)$ is separable, with the notation of Theorem 2.8, we clearly have (recall Proposition 2.6 (i)) $\delta_n \leq \sup\{d_{SC^*(F_\infty)}(E) \mid E \in OS_n\}$, hence the left side of (4.13) follows from (4.12)'''. For the other side, note that by (4.4) and (4.5) we have for any E in OS_n , $d_{SC^*(F_\infty)}(E) \leq d_{SK}(E)$ and by [P3, Theorem 9.6] this is $\leq \sqrt{n}$. ■

Theorem 4.4. *Let A be a C^* -algebra and let $H = \ell_2$. The following are equivalent.*

(i)

$$A \otimes_{\min} B(H) = A \otimes_M B(H).$$

(ii) *For any $\varepsilon > 0$ and any finite dimensional subspace $E \subset A$, there is a subspace $\hat{E} \subset C^*(F_\infty)$ such that $d_{cb}(E, \hat{E}) < 1 + \varepsilon$.*

(iii) *Same as (ii) with $\varepsilon = 0$. Equivalently, every finite dimensional subspace of A is completely isometric to a subspace of $C^*(F_\infty)$.*

Proof. Assume (i). Then for any finite dimensional subspace $E \subset A$, consider a completely isometric embedding $E^* \subset B(H)$ and view $E \otimes E^*$ as a subspace of $A \otimes_{\min} B(H)$. If we apply Proposition 4.2 when $u \in E \otimes E^*$ represents the identity on E , we have $\|u\|_{cb} = 1$, hence $\|u\|_M = 1$ and we immediately obtain (ii). Conversely, (ii) clearly implies (i) by Proposition 4.2 and (4.8). The fact that (i) implies (iii) follows from [EH, Theorem 3.2] (this was kindly pointed out to us by Kirchberg). Indeed, it suffices to prove (iii) for finite dimensional operator systems $E \subset A$. Then, assuming (i) the second condition in [EH, Theorem 3.2] must hold by Lemma 0.7 and the short exactness property of the max-tensor product. Therefore, if we represent A as a quotient of $C^*(F_I)$ for some free group F_I , by [EH, Theorem 3.2] any unital completely positive map $v: E \rightarrow A$ has a unital completely positive lifting $\hat{v}: E \rightarrow C^*(F_I)$. In particular E embeds completely isometrically into $C^*(F_I)$, hence into $C^*(F_\infty)$ (see the proof of Corollary 3.1). This shows (i) $\Rightarrow$ (iii). Finally, (iii) $\Rightarrow$ (ii) is trivial. ■

Remark. The notion appearing in (ii) above is analogous to that of “finite representability” in Banach space theory.

Theorem 4.5. *Let X be an operator space and let $c \geq 1$ be a constant. The following are equivalent.*

- (i) *For any operator space F , we have $X \otimes_{\min} F = X \otimes_M F$ and $\|u\|_M \leq c \|u\|_{\min}$ for any u in $X \otimes F$.*
- (ii) *The same as (i) with $F = B(\ell_2)$.*
- (iii) *For any finite dimensional subspace $E \subset X$ we have*

$$d_{SC^*(F_\infty)}(E) \leq c.$$

Proof. (i) $\Rightarrow$ (ii) is trivial and (ii) $\Rightarrow$ (iii) is clear by Proposition 4.2, taking $F = E^*$ and $u \in X \otimes E^*$ associated to the inclusion $E \subset X$. Finally assume (iii). Consider u in $X \otimes F$. We have $u \in E \otimes F$ for some finite dimensional subspace $E \subset X$. By (iii), for each $\varepsilon > 0$, there is a subspace $\tilde{E} \subset C^*(F_\infty)$ such that $d_{cb}(E, \tilde{E}) \leq c + \varepsilon$. By Proposition 4.2, this implies $\|u\|_M \leq (c + \varepsilon)\|u\|_{\min}$, whence (i). ■

Remark. It can be shown that the operator Hilbert space OH introduced in [P3] satisfies the equivalent conditions in Theorem 4.5 for $c = 1$. In particular we have

$$d_{SC^*(F_\infty)}(OH) = 1.$$

This is a consequence of some unpublished work by U. Haagerup, namely the inequality (4.14) below (itself a consequence of [P3, Corollary 2.7]). To explain this inequality, let $\overline{B(H)}$ be the complex conjugate of $B(H)$, *i.e.* the same space but with the conjugate complex multiplication. We denote by $x \rightarrow \bar{x}$ the canonical anti-isomorphism between $B(H)$ and $\overline{B(H)}$. Note that $\overline{B(H)} \approx B(\overline{H}) \approx B(H)$. In the sequel, we simply denote by $\|\cdot\|_{\max}$ (resp. $\|\cdot\|_{\min}$) the max-norm (resp. the min-norm) on the space $B(H) \otimes \overline{B(H)}$.

Then, for any $x_1, \dots, x_n$ and $y_1, \dots, y_n$ in $B(H)$ we have

$$(4.14) \quad \left\| \sum x_i \otimes \bar{y}_i \right\|_{\max} \leq \left\| \sum x_i \otimes \bar{x}_i \right\|_{\min}^{1/2} \left\| \sum y_i \otimes \bar{y}_i \right\|_{\min}^{1/2}.$$

To check (4.14), we first recall an entirely elementary fact: for any $a_1, \dots, a_n, b_1, \dots, b_n$ in a C^* -algebra A , we have

$$\left\| \sum a_i b_i \right\| \leq \left\| \sum a_i a_i^* \right\|^{1/2} \left\| \sum b_i^* b_i \right\|^{1/2}.$$

Hence if $a_i b_i = b_i a_i$ we also have

$$\left\| \sum a_i b_i \right\| \leq \left\| \sum a_i^* a_i \right\|^{1/2} \left\| \sum b_i b_i^* \right\|^{1/2}.$$

Applying these inequalities in the case $a_i = x_i \otimes 1$, $b_i = 1 \otimes \bar{y}_i$ we get

$$(4.14.0) \quad \left\| \sum x_i \otimes \bar{y}_i \right\|_{\max} \leq \left\| \sum x_i x_i^* \right\|^{1/2} \left\| \sum y_i^* y_i \right\|^{1/2}$$

$$(4.14.1) \quad \left\| \sum x_i \otimes \bar{y}_i \right\|_{\max} \leq \left\| \sum x_i^* x_i \right\|^{1/2} \left\| \sum y_i y_i^* \right\|^{1/2}.$$

Let us denote by A_0 (resp. A_1) the space $B(H)^n$ equipped with the norm $\|(x_i)\| = \left\| \sum x_i^* x_i \right\|^{1/2}$ (resp. $\left\| \sum x_i x_i^* \right\|^{1/2}$). Moreover for any (x_i) in $B(H)^n$, we denote by $\|(x_i)\|_{\frac{1}{2}}$ the norm in the complex interpolation space $(A_0, A_1)_{\frac{1}{2}}$. Then, by the complex interpolation theorem applied to the sesquilinear map

$$(x_i), (y_i) \rightarrow \sum x_i \otimes \bar{y}_i,$$

(4.14.0) and (4.14.1) imply that we have

$$(4.15) \quad \left\| \sum x_i \otimes \bar{y}_i \right\|_{\max} \leq \|(x_i)\|_{\frac{1}{2}} \|(y_i)\|_{\frac{1}{2}}.$$

On the other hand, by Corollary 2.7 in [P3] we have for all (x_i) in $B(H)^n$

$$(4.16) \quad \|(x_i)\|_{\frac{1}{2}} = \left\| \sum x_i \otimes \bar{x}_i \right\|_{\min}^{1/2}.$$

Hence (4.15) implies (4.14).

Finally, let $(T_i)_{i \geq 1}$ be an orthonormal basis in the operator Hilbert space OH introduced in [P3]. We may assume $OH \subset B(H)$ with (say) $H = \ell_2$. Recall (see [P3]) that $\left\| \sum_1^n T_i \otimes \bar{T}_i \right\|_{\min} = 1$ and for any $x_1, \dots, x_n$ in $B(H)$ we have

$$\left\| \sum T_i \otimes x_i \right\|_{\min} = \left\| \sum x_i \otimes \bar{x}_i \right\|_{\min}^{1/2}.$$

Therefore, by (4.14) for any $x_1, \dots, x_n$ in $B(H)$ we have

$$\left\| \sum T_i \otimes x_i \right\|_{\max} \leq \left\| \sum x_i \otimes \bar{x}_i \right\|_{\min}^{1/2} = \left\| \sum T_i \otimes x_i \right\|_{\min}.$$

Equivalently, we conclude that $\|\cdot\|_{\max}$ and $\|\cdot\|_{\min}$ coincide on $OH \otimes B(H)$ so that $X = OH$ satisfies the equivalent properties in Theorem 4.5. Note that if $x_i = y_i$ in (4.14) we have

$$\left\| \sum x_i \otimes \bar{x}_i \right\|_{\max} \leq \left\| \sum x_i \otimes \bar{x}_i \right\|_{\min}.$$

In other words, we obtain that $\|\cdot\|_{\max}$ and $\|\cdot\|_{\min}$ coincide on the “positive” cone in $B(H) \otimes \overline{B(H)}$ formed of all the tensors of the form $\sum_1^n x_i \otimes \bar{x}_i$.

In [P5], a modified version of the identity (4.16) is proved with an arbitrary semi-finite von Neumann algebra in the place of $B(H)$. We refer the reader to a possibly forthcoming paper by U. Haagerup and the second author for extended versions of (4.14) and (4.15).

References

- [AO] C. Akeman and P. Ostrand. Computing norms in group C^* -algebras. Amer. J. Math. 98 (1976), 1015-1047.
- [Ba] R. Baire. Sur les fonctions des variables réelles. Ann. di Mat. (3) 3 (1899) 1-123.
- [B1] D. Blecher. Tensor products of operator spaces II. (Preprint) 1990. Canadian J. Math. 44 (1992) 75-90.
- [B2] _____. The standard dual of an operator space. Pacific J. Math. 153 (1992) 15-30.
- [B3] _____. Tracially completely bounded multilinear maps on C^* -algebras. Journal of the London Mathematical Society, 39 (1989) 514-524.
- [BP] D. Blecher and V. Paulsen. Tensor products of operator spaces. J. Funct. Anal. 99 (1991) 262-292.
- [DCH] J. de Cannière and U. Haagerup. Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups. Amer. J. Math. 107 (1985), 455-500.
- [DHV] P. de la Harpe and A. Valette. La propriété T de Kazhdan pour les groupes localement compacts. Astérisque, Soc. Math. France 175 (1989).
- [EH] E. Effros and U. Haagerup. Lifting problems and local reflexivity for C^* -algebras. Duke Math. J. 52 (1985) 103-128.
- [ER1] _____. A new approach to operator spaces. Canadian Math. Bull. 34 (1991) 329-337.
- [ER2] _____. On the abstract characterization of operator spaces. Proc. Amer. Math. Soc. 119 (1993) 579-584.
- [H1] U. Haagerup. The Grothendieck inequality for bilinear forms on C^* -algebras. Advances in Math. 56 (1985) 93-116.
- [H2] _____. An example of a non-nuclear C^* -algebra which has the metric approximation property. Invent. Math. 50 (1979) 279-293.
- [H3] _____. Injectivity and decomposition of completely bounded maps in "Operator algebras and their connection with Topology and Ergodic Theory". Springer Lecture Notes in Math. 1132 (1985) 170-222.
- [HP] U. Haagerup and G. Pisier. Bounded linear operators between C^* -algebras. Duke

- Math. J. 71 (1993) 889-925.
- [I] T. Itoh. On the completely bounded maps of a C^* -algebra to its dual space. Bull. London math. Soc. 19 (1987) 546-550.
- [Ki1] E. Kirchberg. On subalgebras of the CAR-algebra. (Preprint, Heidelberg, 1990) J. Funct. Anal. (To appear)
- [Ki2] _____. On non-semisplit extensions, tensor products and exactness of group C^* -algebras. Invent. Math. 112 (1993) 449-489.
- [Ki3] _____. Commutants of unitaries in UHF algebras and functorial properties of exactness. J. reine angew. Math. To appear.
- [Kr] J. Kraus. The slice map problem and approximation properties. J. Funct. Anal. 102 (1991) 116-155.
- [Ku] W. Kuratowski. Topology. Vol. 1. (New edition translated from the french.) Academic Press, New-York 1966.
- [Kw] S. Kwapien. On operators factorizable through L_p -spaces. Bull. Soc. Math. France, Mémoire 31-32,(1972) 215-225.
- [La] C. Lance. On nuclear C^* -algebras. J. Funct. Anal. 12 (1973) 157-176.
- [Pa1] V. Paulsen. Completely bounded maps and dilations. Pitman Research Notes 146. Pitman Longman (Wiley) 1986.
- [Pa2] _____. Representation of Function algebras, Abstract operator spaces and Banach space Geometry. J. Funct. Anal. 109 (1992) 113-129.
- [Pa3] _____. The maximal operator space of a normed space. To appear.
- [P1] G. Pisier. Exact operator spaces. Colloque sur les algèbres d'opérateurs (Orléans 1992). Astérisque. Soc. Math. France. To appear.
- [P2] _____. Factorization of linear operators and the Geometry of Banach spaces. CBMS (Regional conferences of the A.M.S.) 60, (1986), Reprinted with corrections 1987.
- [P3] _____. The operator Hilbert space OH , complex interpolation and tensor norms. Memoirs Amer. Math. Soc. (submitted)
- [P4] _____. Factorization of operator valued analytic functions. Advances in Math. 93 (1992) 61-125.

- [P5] _____. Projections from a von Neumann algebra onto a subalgebra.
Bull. Soc. Math. France. To appear.
- [Ru] Z. J. Ruan. Subspaces of C^* -algebras. J. Funct. Anal. 76 (1988) 217-230.
- [Sa] S. Sakai. C^* -algebras and W^* -algebras. Springer Verlag New-York, 1971.
- [Sm] R. R. Smith. Completely bounded maps between C^* -algebras. J. London Math. Soc. 27 (1983) 157-166.
- [Ta] M. Takesaki. Theory of Operator Algebras I. Springer-Verlag New-York 1979.
- [Tr] S. Trott. A pair of generators for the unimodular group. Canad. Math. Bull. 3 (1962) 245-252.
- [V] D. Voiculescu. Property T and approximation of operators. Bull. London Math. Soc. 22 (1990) 25-30.
- [VDN] D. Voiculescu, K. Dykema, A. Nica. Free random variables. CRM Monograph Series, Vol. 1, Amer. Math. Soc., Providence RI.
- [W1] S. Wassermann. On tensor products of certain group C^* -algebras. J. Funct. Anal. 23 (1976) 239-254.
- [W2] _____. Exact C^* -algebras and related topics. Lecture notes series 19, Seoul National University, 1994.

M. Junge:
 Mathematisches Seminar
 CAU Kiel,
 24098 Kiel, Germany

G. Pisier:
 Texas A&M University
 College Station, TX 77843, U. S. A.
 and
 Université Paris 6
 Equipe d'Analyse, Boîte 186,
 75252 Paris Cedex 05, France